

PRODUCT INFORMATION

WSD KEYMIN

For Cattle and Sheep

Product Description

WSD KEYMIN is a broad spectrum mineral supplement for ruminant livestock and is an aid in managing selenium, cobalt or copper deficiencies.

Type of Preparation

WSD KEYMIN is a solution to be added to water or administered orally containing chelated mineral nutrients.

Indications

To correct imbalances or deficiencies of minerals in cattle, sheep and goats. The use of chelated minerals ensures their availability to the animal when administered orally. A high quality supplement to be administered on an intermittent basis in water or dosed orally when:

- a deficiency of one or more minerals might be expected or
- to aid in digestion by rumen microbes when fodder low in one or more inorganic nutrients is being consumed by sheep, cattle or goats.

Milking Sheep, Dairy Cattle & Dairy Goats: Suitable for use in animals producing milk for human consumption.

Dosage

Always refer to the label. The dose rate is according to body weight range. This product may be added to drinking water so that animals are dosed at the label rate or is suitable for injection into stock water supplies by metered devices.

If mixed with water, it is preferable to administer the required dose over 2 to 3 days until the recommended total average dose is received by each individual animal.

Further doses may be indicated at intervals of 2 to 4 weeks during the period when supplementation is required.

Advice On Use

- **WSD KEYMIN** should be administered undiluted when drenching sheep, cattle or goats.
- **WSD KEYMIN** is used to supplement the diet of ruminants, particularly on mature dry feed, when moderate to high levels of grain are being fed, or when grass pastures are growing rapidly.
- This product should not be administered at a rate greater than recommended on the label.
- If this product is added to water there is no need to train or adapt animals to consuming **WSD KEYMIN**.
- It is recommended that **WSD KEYMIN** should not be mixed with other oral drenches. Do NOT mix with preparations containing levamisole.
- Goats may be dosed at the same rate as sheep.
- Care should be exercised if copper bullets have been administered or if livestock are grazing plants likely to cause liver damage.
- WSD produces a range of mineral supplements for different purposes and situations including **WSD KEYMIN**, **WSD FACTOR G** and **WSD HYTONE**.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



Keymin

Active constituent:

20 mg/L Chromium as Lignosulfonate
6.40 g/L Copper as Lignosulfonate
1.10 g/L Cobalt as Lignosulfonate
8.50 g/L Zinc as Lignosulfonate
3.90 g/L Magnesium as Lignosulfonate
0.03 g/L Iron as Lignosulfonate
6.20 g/L Manganese as Lignosulfonate
0.03 g/L Boron as Lignosulfonate

3.20 g/L Phosphorus as Lignosulfonate
1.00 g/L Selenium as Selenate/Selenite
1.60 g/L Calcium as Lignosulfonate
12.2 g/L Sodium as Lignosulfonate
42.5 g/L Sulfur as Lignosulfonate
200 g/L Crude protein
65.7 g/L Glucose

Chelated trace element supplement for sheep and cattle
to supplement diets where mineral levels may be low.

20 L



DIRECTIONS FOR USE:

Make sure contents are well mixed by shaking container prior to use.
Ensure correct drenching rates are followed and avoid injury to livestock by maintaining a smooth drenching nozzle.

NO WITHHOLDING PERIOD APPLIES WHEN USING THIS PRODUCT ALONE.

WARNING: THIS PRODUCT MUST NOT BE APPLIED AT A RATE GREATER THAN THE AMOUNT SHOWN ON THE LABEL. NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

AS A DIARY AND BEEF CATTLE DRENCH			
Liveweight (kg)	Drench Rate	Number of Cattles per Pack	Minimum Frequency (weeks)
0-50	5mL	4000	2
50-100	10mL	2000	2
101-150	15mL	1333	3
151-200	20mL	1000	3
201-250	25mL	800	4
251-300	30mL	666	4
301-350	35mL	571	5
351-400	40mL	500	6
401-450	45mL	444	6
451+	50mL	400	6

AS A FEED OR WATER ADDITIVE FOR DAIRY AND BEEF CATTLE		
Liveweight (kg)	Daily Rate per Animal	Rate per Animal (weekly)
0-50	0.50mL	3.50mL
50-100	1.50mL	10.50mL
101-150	1.50mL	10.50mL
151-200	2.00mL	14.00mL
201-250	2.00mL	14.00mL
251-300	2.00mL	14.00mL
301-350	2.00mL	14.00mL
351-400	2.00mL	14.00mL
401-450	2.00mL	14.00mL
451+	2.50mL	17.50mL

AS A DRENCH FOR SHEEP			
Liveweight (kg)	Drench Rate	Number of Sheep per Pack	Minimum Frequency (weeks)
0-30	5mL	4000	6
31-40	6mL	3333	6
41-50	8mL	2500	6
51-70	10mL	2000	6
70+	12mL	1666	6

AS A FEED OR WATER ADDITIVE FOR SHEEP		
Liveweight (kg)	Daily Rate per Animal	Rate per Animal (weekly)
15-19	0.15mL	1.00mL
20-25	0.20mL	1.50mL
26-30	0.25mL	1.75mL
31-40	0.30mL	2.00mL
41-50	0.40mL	3.00mL
51-70	0.50mL	3.50mL
70+	0.55mL	4.00mL

SAFETY DIRECTIONS:

Poisonous if absorbed by skin contact or swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container or using the product wear rubber gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating and drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone in Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

DISPOSAL:

Triple or preferably pressure rinse. Do not dispose of undiluted chemicals on-site. Break, crush or puncture and bury containers in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

STORAGE:

Store below 30°C (Room Temperature) in original container in a dry place. Protect from light. Keep container well closed.

CONDITIONS OF SALE:

All WSD Agribusiness products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

Manufactured and distributed in Australia by:



WSD Agribusiness Pty. Ltd.
Koojan Avenue, South Guildford WA 6055
Tel: (08) 9321 2888 Fax: (08) 9479 4088
contact@wsdagribusiness.com
www.wsdagribusiness.com

Batch:

Expiry:



9 330805 001133